

B

Ullmann's Encyclopedia of Industrial Chemistry

Fifth, Completely Revised Edition

Volume A 5:

Cancer Chemotherapy to Ceramic Colorants

Executive Editor: Wolfgang Gerhartz

Senior Editor: Y. Stephen Yamamoto

Editors: F. Thomas Campbell, Rudolf Pfefferkorn,
James F. Rounsaville

Henkel KGaA
Chem. Bldg.



BEST AVAILABLE COPY

1986

Vol. A5

Carbohydrates 79

Carbohydrates

The following are separate keywords: Cellulose, Cellulose Esters, Cellulose Ethers, Cellulose Fibers, Fructose, Gluconic Acid, Glucose and Glucose-Containing Symp., Lactose, Polysaccharides, Sorbitol, Starch and Starch Derivatives, Sugar, Sugar Alcohol

JOCHEN LIEBMAN, Institut für Organische Chemie und Biochemie, Universität Freiburg, Freiburg, Federal Republic of Germany
 KNUST RAFT, Südmeier AG, Gießen, H-Oberhessen, Federal Republic of Germany

1. Monosaccharides	80	5. Reactions of Carbohydrates	85
2. Oligosaccharides	83	5.1. Reactions of the Hydroxyl Groups	87
3. Polysaccharides	83	5.2. Reactions of the Carbonyl Group	90
4. Nomenclature	84	6. References	92

The term "carbohydrates" describes a major group of naturally occurring compounds of widely varying molecular masses. The low molecular mass carbohydrates comprise a family of more or less sweet-tasting, water-soluble, colorless products with physical, chemical, and physiological properties. They are often generically called "sugars," which is the familiar sucrose.

The high molecular mass carbohydrates are called polysaccharides. Their physical properties differ significantly from those of low molecular mass. They do not taste sweet and many do not dissolve in water.

All carbohydrates are composed of units of monosaccharides or their derivatives. Carbohydrates are usually classified by the number of monosaccharide units in their molecules. A monosaccharide contains an uninterrupted chain of a limited number (five or six) of carbon atoms. In disaccharides, trisaccharides, tetrasaccharides, etc., and in polysaccharides, n monosaccharides are linked together through glycosidic oxygen atoms.

Oligosaccharides are considered to be low molecular mass carbohydrates. However, no definite borderline between oligosaccharides and the polysaccharides exists. In general, naturally occurring oligosaccha-

rides a major chemical component of many biological systems. Carbohydrates comprise a family of more or less sweet-tasting, water-soluble, colorless products with physical, chemical, and physiological properties. They are often generically called "sugars," which is the familiar sucrose.

Most monosaccharides, oligosaccharides, and polysaccharides have the common molecular formula $C_n(H_2O)_m$ (from which the common name "carbohydrate" is derived), in which n is either equal to m (monosaccharides) or a little larger than m (oligo- and polysaccharides). Oligomers are formed by condensation (formal elimination of water) of monosaccharide units, and conversely oligo- and polysaccharides can be degraded to monosaccharides by hydrolysis. The whole biomass on earth has been estimated to be about 90% carbohydrates, which means that the molecular formula of the total organic matter on earth is only slightly different from that of the carbohydrates.

Carbohydrates have important functions: primarily they supply chemical energy to the living cell and serve as raw materials for the construction of other natural products. Carbohydrates are mainly formed in the photochemical carbon dioxide fixation in green plants, a complicated biochemical process which provides for the efficient conversion of solar energy into chemical energy. The uses of carbohydrates for the supply of primary energy and as raw materials in synthetic organic chemistry represent an important facet of present and future technology.

Carbohydrates are industrially utilized to a considerable extent. Glucose [50-99-7] is the

starting material for the product (ascorbic acid). Maltose [69-7] is the enzymatic degradation of duced to maltitol, which is use sweetener. A similar sweetener nh) [64519-83-0], a substance o alcohols *O*- α -D-glucopyranoside (1) [20942-99-8] and *O*- α - (1 \rightarrow 6)-*D*-glucinol [334-73-6], tained by reduction of isomalt [13718-94-0], a product of enation of sucrose. Fructose [57-4, a sweetener by diabetics.

Many nonionic detergent carbohydrates. Used in soap are less irritating than their acid counterparts and can easily be degraded before causing too environmental problems. Some are produced and used in a variety of applications. These include: blood serum (dextran), and starch (amylose), wrapping films, additives to food and pharmaceuticals and cosmetics (starches, plant and microbial oils), fluids in oil fields. Cellulose is or cotton is one of the most important raw materials.

Production. Natural cart tained exclusively from biolo; synthoses are impractical, as 3-epimerization of D-glucose; saccharide, D-allose, which ture, is tedious. Chemical saccharides by coupling suit monosaccharides are extre only possible on a small scale thesis of a polysaccharide is One possibility is the poly anhydro sugar derivatives, levoglucosans, which yields pyranans.

1. Monosaccharides

Monoaccharides are either aldehydes (-oses, aldoses) or p (-uloses, ketoses). The carbon monosaccharides is linear (in most all known the carbonyl carbon atom 2. Most monosaccharides are hexoses and hexuloses).

on of vitamin C
6), a product of
arch, can be re-
as a low-calorie
isomalt (palati-
the disaccharide
-(1 → 1)-D-man-
glucopyranosyl-
which are ob-
loose (palatinose)
matic isomeriza-
7) itself is used as


are derived from
r shampoo, they
nic counterparts
ologically, there-
problem. Poly-
sed in a wide va-
clude substitutes
esives (cellulose
material (collo-
beverages, phar-
m, pectic sub-
), and drilling
be form of wood
ortant industrial

hydrates are ob-
al sources. Total
even the simple
to another mono-
not found in na-
theses of oligo-
als derivatives of
ly difficult and
the chemical syn-
most impossible.
erization of 1,6-
ch as substituted
1 → 6)-α-D-gluc-

polyhydroxy aldehydes (polyhydroxy ketones skeleton in most branched). In aldehydes, the aldehyde group is located at the end of the chain. Monosaccharides have either three (trioses) or four (tetroses) or five (pentoses)

and pentuloses) carbon atoms. Monosaccharides with fewer carbon atoms (trioses, tetroses, and tetrols) or more carbon atoms (heptoses, octoses, heptuloses, octuloses, etc.) rare. The most important and abundant monosaccharides are hexoses: D-glucose, D-mannose, and D-galactose. D-fructose is a hexulose. Ubiquitous pentoses are L-arabinose, D-xylose, and D-ribose.

Stereoisomerism. Monosaccharides (as polyhydroxy aldehydes or ketones) possess several asymmetric carbon atoms. Hexoses have four, hexuloses and pentoses three, and pentuloses two chirality centers. The existence of several asymmetric carbon atoms in one molecule gives rise to families of stereoisomers. For a hexose with four asymmetric carbon atoms, there exist 2^4 stereoisomers, eight of which are enantiomers (mirror images) of the other eight. Hexuloses exist in four pairs of enantiomers, pentuloses and tetroses in pairs. The diastereomeric monosaccharides (diastereomers are stereoisomers that are not mirror images of each other) have trivial names. In Figure 1, D-hexoses, D-hexuloses, D-pentoses, and D-tetroses are represented as open-chain Fischer projections. In the Fischer projection all carbon atoms form a bow with the convex part pointing toward the viewer. According to the Fischer convention, the orientation of the hydroxyl group attached to the highest numbered asymmetric carbon atom determines whether a monosaccharide belongs to the D- or the L-series. If this group is positioned on the right of the carbon chain, assignment is to the D-series, and vice versa. Most naturally occurring monosaccharides have the D-configuration. The chirality of carbohydrates makes this class of natural compounds an ideal source for the preparation of chiral synthons (intermediates for chemical synthesis), which are needed for the synthesis of pharmaceuticals, agrochemicals, pheromones, etc. [26], [27].

**Figure 1**

Pyranoses and Furanoses. Monosaccharides form cyclic intramolecular hemiacetals if their carbon skeleton permits. The relative stabilities of these hemiacetals depend on the ring size. In general, six-membered rings (pyranoses) are most common, whereas only a few five-membered rings (furanoses) are stable. Smaller rings are unstable, and larger ones can only be detected in aqueous equilibrium mixtures. Structures of the cyclic hemiacetals are represented by the Haworth projection formulas of the D-series



Figure 2.
Verdict

02-JUN 18:42 FR UON:WME ED SMULDERS

+49 211 7988688

RN: +49 211 79814011

SEITE: 04

Vol. A5

Cellulose 375

Cellulose

HANS KILBING, Seewalchen, Austria, Josef SCHURZ, University of Graz, Austria (Chap. 1)
 ROBERT G. STRADMAN, Texas Tech University, Lubbock, Texas, United States (Chap. 2)
 KARL SCHLUNGER, Textilforschungsanstalt Krefeld, Krefeld, Federal Republic of Germany (Chap. 2.1.1)
 WILHELM ALBRECHT, Wuppertal Federal Republic of Germany (Chap. 3)

1. Cellulose	375	2.3. Jute	398
1.1. Properties	376	2.4. Flax	399
1.1.1. Molecular Structure	376	2.5. Hemp	399
1.1.2. Supermolecular Structure (Texture)	380	2.6. Leaf (Hard) Fibers	399
1.1.3. Physical Properties	383	2.7. Cane	400
1.1.4. Chemical Properties	383	2.8. Economic Aspects	400
1.2. Occurrence	389	2.9. Occupational Health	400
1.3. Production	390	3. Regenerated Cellulose	400
1.4. Quality Testing	390	3.1. Viscose Fibers	401
1.5. Applications	391	3.1.1. Principles of the Viscose Process	401
2. Natural Cellulose Fibers	391	3.1.2. Viscose Preparation	402
2.1. Cotton	392	3.1.3. Viscose Fiber Spinning	404
2.1.1. Molecular Arrangement, Morphology, and Fine Structure	392	3.1.4. Fiber Types	406
2.1.2. Properties	394	3.1.5. Modified Viscose Fibers	408
2.1.3. Production	396	3.1.6. Fiber Properties	409
2.1.3.1. Harvesting	396	3.1.7. Uses	412
2.1.3.2. Ginning	396	3.1.8. Economic Aspects	412
2.1.3.3. Byproducts	396	3.2. Cuprammonium Fibers	413
2.1.3.4. Processing	397	3.3. Tentative Cellulose Fiber Production by Other Processes - Outlook	415
2.1.3.5. Finishing	397	4. References	416
2.1.3.6. Special Finishes	397		
2.2. Bast (Soft) Fibers	398		

1. Cellulose

Cellulose [9004-34-6] deserves a special position among the industrially used raw materials for two general reasons. First, as to the natural products which, fully, are inexhaustible since it is generated by nature in relatively short time periods. As long as we ensure that the per cellulose, forests and cotton plant damaged by destructive lumbering, we can expect regular and significant annual reproduction.

According to references [8], if cellulose matter resulting from

a special position among the industrially used raw materials for two general reasons. First, as to the natural products which, fully, are inexhaustible since it is generated by nature in relatively short time periods. As long as we ensure that the per cellulose, forests and cotton plant damaged by destructive lumbering, we can expect regular and significant annual reproduction.

biosynthesis amounts to approximately 1.3×10^9 metric tons. A tree produces an average of 13.7 g of cellulose daily. If they were lined up, the cellulose chain molecules formed each day would result in a string of 2.62×10^{10} km in length, or 175 times the distance between the sun and the earth.

In wood, cellulose is part of an ingeniously constructed fiber-reinforced composite in which long, stiff cellulose chain molecules organized in thin fibrils constitute the plant skeleton material held together and protected by hydrophobic lignin acting as binder and encasement.

To isolate cellulose from wood for industrial applications, the wooden composite must be broken up by so-called pulping processes. In

02-JUN 10:43 FR UON:MAE ED SHILDERS

+49 211 7986680

ANI+49 211 79814811

SEITE:05

876 Cellulose

these treatments, other wo as lignin and hemicellulose degraded and dissolved. T products have found only limit wood pulp manufacturers pulping liquors to concent 50% solids. The organic n produce steam and electric ganic pulping chemicals ammonium base and sulfi nceously recovered. Ther have practically solved t rontmental problems of t

Both cellulose and lig gradable and, thus, coolo will decompose in the op such as paper or cellulor pose and eventually form dustrial use, environme caused by cellulose or lig used in the isolation or processing and transfo derivativum, films, or fibe term task of modern cell development of novel p or only a few ecologically these efforts are succes maintain and strengthen able and environmental important raw material cally produced polymer


constituents, such are to a large extent s far, these byprod . use. In most cases, monstrate the waste tes consisting of ca. lter is used as fuel to over, while the inor- xida, magnesium, or dioxide) are simulta- recovery processes long-standing envi- wood pulp industry. s are biologically de- ally beneficial. They . Cellulose products, textiles, will decomp- aturable humus. In in- al problems are not t but by the chemicals subsequent chemical uation into cellulose . Therefore, the long- es research will be the cases which yield no arnful byproducts. If l, cellulose will surely a position as a recow- beneficial, industrially mpeting with syntheti-

Cellulose products, textiles, will decomp- aturable humus. In in- al problems are not t but by the chemicals subsequent chemical uation into cellulose . Therefore, the long- es research will be the cases which yield no arnful byproducts. If l, cellulose will surely a position as a recow- beneficial, industrially mpeting with syntheti-

glucose base unit is $m_0 = 162$, and the molecular mass of the cellulose polymer is

$$M_n = m_0 P + 18 = 162 P.$$

Constitutional Formula. HAWORTH [10] first discovered the covalent bonds inside and between the glucose units while STAUDINGER [11] found the final proof for the macromolecular nature of the cellulose molecule.



Conformational Formula. The glucopyranose ring adopts a 4_1 chair conformation, as revealed by recent X-ray crystallography and nuclear magnetic resonance studies [12], [13] with glucose. The chair formation in comparison to the tray conformation exhibits a free stabilization enthalpy of $G_s = 20.05$ kJ/mol [14]. In this conformation, the three hydroxyl groups are positioned in the ring plane while the hydrogen atoms are in a vertical position. It seems only natural to assume that the same conformation also exists in the cellulose molecule.



1.1. Properties

1.1.1. Molecular Structure

Cellulose is an insoluble β -1,4-polyacetal of cellobiose (4-O- β -D-glucopyranosyl-D-glucose). The actual base unit, cellobiose, consists of two molecules of glucose can also be considered as a polyacetal of glucose.

Basic Structure. The basic chemical formula of cellulose is the following:




where P = the degree of polymerization; n = the number of units in the chain.

The elemental composition of cellulose was already known to PAYEN in 1842 [9]. The

basic chemical formula of cellulose is the following:

position of 44.4% C, 6.2% H, and 49.4% O was already known to PAYEN in 1842 [9]. The molecular mass of the



Structural Anomalies. As a naturally occurring polymer, cellulose always contains small amounts of other constituents in addition to glucose (over 99%). These may already be partially built into or onto the cellulose molecules during biosynthesis, such as lignin-cellulose complexes [15]. Most of the changes in the molecular structure, however, result from secondary reactions, i.e., hydrolysis or oxidation, during isolation from natural sources. For morphological reasons, such chemical changes occur preferably in

SEITE:06

Type of cellulose	η_{sp}
Cotton, raw	7 000
Cotton, raw (according to Russian spec)	14 000
Cotton, purified	1 500-300
Cotton linters	6 500
Flax	8 000
Ramie	6 200
α -Cellulose (isolated from wood fibers)	1 100-800
Spruce, pulped	3 300
Beech, pulped	3 050
Aspen	2 500
Fir	2 500
Bacterial cellulose	2 700
Acetobacter cellulose	600

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.